



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Equation 2, $x_2^2 - px_2 + q = 0$, may be solved in a similar manner by changing the sign of px and proceeding as above. The roots being the same in numerical value but opposite in sign.


Case II. Equations 3, 4. The figure for Case II differs from that of Case I only in that P is outside of the circle 1; the points CPD being now in the order PCD . $PC = x_1$, $PD = x_2 = x_1 + p$, $PN \cdot PM = PC \cdot PD$; $r \cdot s = x_1(x_1 + p)$, $x_1^2 + px_1 - rs = 0$. Similarly, $x_2^2 + px_2 - rs = 0$.

Equation 4 is solved by changing sign of px and proceeding as with equation 3, since the roots are the opposite of the roots of equation (3).

In Case I if the roots are imaginary, the point P falls within circumference 2, and the graphic method fails.

VI. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Upon AB describe a semi-circle. Let C be the mid-point of the semi-circular arc. Draw AC , BC . Let G be a point on the line AC ; on GC describe a circle center D . Through D draw $BFDE$. Then taking BE , BF positive; EB , FB negative, these lines are the roots of a quadratic having $2CD$ for the coefficient of the first power of the unknown quantity, and $\sqrt{2AB}$ for the absolute term, the coefficient of the second power of the unknown quantity being taken unity.



Let $AB = \sqrt{2}c$, $GC = 2DC = b$.

$BE = BD + DE = BD + DC = \sqrt{BC^2 + DC^2} + DC = \frac{1}{2}[b + \sqrt{(b^2 + 4c)}]$.

$BF = BD - DF = BD - DC = -\frac{1}{2}[b - \sqrt{(b^2 + 4c)}]$. Taking $x^2 \pm bx = c$.

then for $x^2 + bx = c$, $EB = -\frac{1}{2}[b + \sqrt{(b^2 + 4c)}]$, $BF = -\frac{1}{2}[b - \sqrt{(b^2 + 4c)}]$.

For $x^2 - bx = c$, $BE = \frac{1}{2}[b + \sqrt{(b^2 + 4c)}]$, $FB = \frac{1}{2}[b - \sqrt{(b^2 + 4c)}]$.

If c be negative, the results still hold.

Also solved by G. W. Greenwood, B. A. (Oxon), Professor of Mathematics and Astronomy in McKendree College, Lebanon, Ill., by use of circle and hyperbola.

219A. Proposed by H. F. MacNEISH, A.B., Assistant in Mathematics, University High School, Chicago, Ill.

Draw a line through a given point which shall divide a given quadrilateral into two equivalent parts; (1) when the point lies in a side of the quadrilateral, (2) when the point is without, (3) within the quadrilateral.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let $ABCD$ be the given quadrilateral. Produce DA , CB till they meet in F . Join AC and draw BS parallel to AC , join CS ; then triangle $SCD =$ quadrilateral $ABCD$. Disect SD in H , HF in G , and join CH , CG .


(1). Let P be the point in the side BC . Join PH and draw CL parallel to PH , join PL . Triangle $PCL =$ triangle HCL .

$\therefore PCL + LCD = PCDL = HCL + LCD = HCD = \frac{1}{2}ABCD$.

(2). Let R be the point without the quadrilateral. Draw RK parallel to

FD meeting FC in K . Join KG and draw CI parallel to KG , join KI . Then $ICG=ICK$ and $FCI=FCI$.

$\therefore FCG=FKI$. Draw IJ parallel to FC , then parallelogram $FKJI=FCH$. At the point I draw IZ perpendicular to FD and equal to RK , draw $ZL=RJ$.



Then $RPUL$ is the line required; for $LZ^2 - IZ^2 = IL^2$ or $VRJ - PRK = IVL = PVJK$.

$\therefore FPL = FKJI = FCH$. Now $SBA = SBC$. $\therefore FBA = FCS$ or $FPL - FBA = FCH - FCS$.

$\therefore ABPL = SCH = \frac{1}{2}ABCD$.

(3). Let Q be the point within the quadrilateral. Draw QE parallel to FD meeting FC in E . Join EG and draw CM parallel to EG , join EM . Then $EGM = EGC$ and $FEM = FCG$. Draw MN parallel to FC ; then $FENM = FCH$. At the point M draw MY per-

pendicular to FD and equal to EQ , draw $YL = QN$. Then $PQOL$ is the required line, for $LY^2 = MY^2 + ML^2$ or $QON = PEQ + MOL$.

$\therefore FPL = FENM = FCH$. $\therefore FPL - FBA = FCH - FCS = SCH$.

$\therefore ABPL = SCH = \frac{1}{2}ABCD$.

Also solved by G. W. Greenwood, B. A. (Oxon).

220. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons, West Va.

Two triangles are circumscribed to a given triangle ABC , having their sides perpendicular to the sides of the given triangle. Prove that the two triangles are equal, and find the area of these triangles.

I. Solution by G. W. GREENWOOD, B. A. (Oxon), Professor of Mathematics and Astronomy, McKendree College, Lebanon, Ill.

The perpendiculars to a side of the given triangle at its extremities, which are corresponding sides of the circumscribed triangles, are symmetrical with respect to any point on the perpendicular bisector of that side. Hence the two triangles are symmetric with respect to the circumcenter of the given triangle and are therefore equal. The area of either is equal to

$$\frac{1}{2}(c^2 \cot B + a^2 \cot C + b^2 \cot A) + S,$$

where S is the area of the given triangle.